


Development of novel Anticancer drugs using *in silico* studies on Aurora Kinase B using PubChem Database



Mrs. Ipsa AjitKumar Singh
(Ph.D. Scholar)

Bioinformatics Research Laboratory

**Dr. D. Y. PATIL BIOTECHNOLOGY AND BIOINFORMATICS
INSTITUTE, TATHAWADE,
Dr. D. Y. PATIL VIDYAPEETH, PUNE**

Introduction and Background


- **Aurora kinases**
 - Group of enzymes belong to the Serine- threonine protein kinase family.
 - It plays an important role in cell division and participates in several steps of mitosis and chromatid segregation .
 - Overexpression, polymorphism and splicing variants in the Aurora Kinase B leads to tumorigenesis and further it leads to cancer .

Location, function and tumor types in different Aurora kinases

Kinases	Localization	Function	Tumour type
Aurora Kinase B	Chromosome Kinetochore Midbody	Chromosome condensation; Microtubule-kinetochore attachment; Chromosomal alignment; Chromosomal segregation; Regulating Cytokinesis	Breast cancer, Ovarian cancer, Gastric/Gastrointestinal cancer, Colorectal cancer, Lung cancer, Cervical cancer, Prostate cancer, Glioma, Acute myeloid leukemia (AML), Oral cancer


Flavones derivatives


- Flavones are a class of flavonoids.
- Common in fruits and many plant foods.
- Shows anticancer activity.
- Flavone derivatives containing fluoro groups are taken.
- These fluoro derivative flavones with Aurora Kinase B is used in docking experiments .




Filtration

Total number of analogs
of Fluoro Flavones from
Pubchem databases =
3882






Using LigPrep



Protein Preparation for docking





- Aurora Kinase B



Aurora Kinase B
PDB ID: 4AF3

Protein is
ready for
docking


Docking results

Glide Docking	PubChem (structure)	ID	2 D Structure	Binding Energy (kcal/mol)	Interaction	Bond Type	Bond length (Å)
SP	44298667		 	-9.153	Pro158	H-bond	1.87
					Ala157	H-bond	2.02
					Ala157	Aromatic H-bond	2.45
					Ala157	Aromatic H-bond	2.54
XP	101664315			-10.287	Ala157	H-bond	2.00
					Tyr156	Aromatic H-bond	3.20
Reference	261400			-8.907	Ala157	H-bond	2.00
					Tyr156	Aromatic H-bond	3.18
					Ala157	Aromatic H-bond	2.63



Interactions with Aurora Kinase A

PubChem ID	CID: 44298667 (SP Docking)	CID:101664315 (XP Docking)	CID: 261400
Binding Energy (kcal/mol)	-9.153	-10.287	-8.907
2D Structure			
Interactions			

Enrichment calculations




Molecular dynamic (MD) studies



Time dependence of root mean square deviations (RMSDs) of the C- α backbone of Aurora Kinase A after binding with fluoro flavones and its 2 best analogs with viz. CID: 261400, 44298667, 101664315

Molecular dynamic (MD) studies (continued)



Time dependence of root mean square fluctuation (RMSFs) of the C- α backbone of Aurora Kinase A after binding with floro flavones and its 2 best analogs with viz. CID: 261400, 44298667, 101664315.

Prime MM/GBSA energies for the complexes of Aurora Kinase A protein

Flouro flavone derivatives (CID)	ΔG Bind ¹ (kcal/mol)	ΔG Coulomb ² (kcal/mol)	ΔG Bind vdW ³ (kcal/mol)	ΔG Solv GB ⁴ (kcal/mol)	Complex Energy ⁵ (kcal/mol)
261400	-56.051 ± 1.942	-2.622 ± 1.001	8.458 ± 1.554	-33.201 ± 1.294	-8114.447 ± 2.0628
101664315	-62.181 ± 1.270	-6.359 ± 0.397	7.384 ± 0.543	-31.059 ± 2.001	-8294.082 ± 1.174
44298667	-65.981 ± 1.995	-16.471 ± 0.644	10.203 ± 1.014	-32.771 ± 1.607	-8148.007 ± 2.074

¹ MM/GBSA binding free energy.

² Coulomb energy.

³ Van der Waals energy.

⁴ GB Generalized Born electrostatic solvation energy.

⁵ Energy of protein-ligand complex

Conclusion

- The result obtained show that 2-(1,1,2,2,3,3-Hexafluoropropyl)-4H-1-benzopyran-4-one has best docking score of -10.287 kcal/mol among all the analogs docked with Aurora Kinase B
- MD simulations was carried out for of 100 ns for each of these three complexes of Aurora Kinase B and flouro flavones and its 2 best analogs, with the mean RMSD value of 2.56Å.
- The molecular dynamics results further confers the conformational stability of the protein-ligand complexes as the RMSF of the pocket residues after binding of the ligand and the protein is lower than 2.0 Å .
- The MM/GBSA free energy calculations reveal that the analogs of the compound forms a stable complex with Aurora Kinase B.
- Thus, the obtained 2 best analogs of flouro flavones can be further used *in vitro* and *in vivo* experiments and can probably serve as an novel drug for cancer treatment.

References

1. Carvalhal, S., Ribeiro, S. A., Arocena, M., Kasciukovic, T., Temme, A., Koehler, K., Huebner, A., & Griffis, E. R. (2015). The nucleoporin ALADIN regulates Aurora A localization to ensure robust mitotic spindle formation. *Molecular biology of the cell*, 26(19), 3424–3438.
<https://doi.org/10.1091/mbc.E15-02-0113>
2. Kamran, M., Long, Z. J., Xu, D., Lv, S. S., Liu, B., Wang, C. L., Xu, J., Lam, E. W., & Liu, Q. (2017). Aurora kinase A regulates Survivin stability through targeting FBXL7 in gastric cancer drug resistance and prognosis. *Oncogenesis*, 6(2), e298. <https://doi.org/10.1038/oncsis.2016.80>
3. Tang, A., Gao, K., Chu, L., Zhang, R., Yang, J., & Zheng, J. (2017). Aurora kinases: novel therapy targets in cancers. *Oncotarget*, 8(14), 23937–23954. <https://doi.org/10.18632/oncotarget.14893>
4. D'Assoro AB, Haddad T, Galanis E. Aurora-A Kinase as a Promising Therapeutic Target in Cancer. *Front Oncol.* 2016 Jan 6;5:295. doi: 10.3389/fonc.2015.00295. PMID: 26779440; PMCID: PMC4701905.
5. Shin, S. Y., Lee, Y., Kim, B. S., Lee, J., Ahn, S., Koh, D., Lim, Y., & Lee, Y. H. (2018). Inhibitory Effect of Synthetic Flavone Derivatives on Pan-Aurora Kinases: Induction of G2/M Cell-Cycle Arrest and Apoptosis in HCT116 Human Colon Cancer Cells. *International journal of molecular sciences*, 19(12), 4086. <https://doi.org/10.3390/ijms19124086>
6. Yan, M., Wang, C., He, B., Yang, M., Tong, M., Long, Z., Liu, B., Peng, F., Xu, L., Zhang, Y., Liang, D., Lei, H., Subrata, S., Kelley, K. W., Lam, E. W., Jin, B., & Liu, Q. (2016). Aurora-A Kinase: A Potent Oncogene and Target for Cancer Therapy. *Medicinal research reviews*, 36(6), 1036–1079. <https://doi.org/10.1002/med.21399>
7. Lee, Y. H., Park, J., Ahn, S., Lee, Y., Lee, J., Shin, S. Y., Koh, D., & Lim, Y. (2019). Design, synthesis, and biological evaluation of polyphenols with 4,6-diphenylpyrimidin-2-amine derivatives for inhibition of Aurora kinase A. *Daru : journal of Faculty of Pharmacy, Tehran University of Medical Sciences*, 27(1), 265–281. <https://doi.org/10.1007/s40199-019-00272-5>
8. Katayama, H., & Sen, S. (2010). Aurora kinase inhibitors as anticancer molecules. *Biochimica et biophysica acta*, 1799(10-12), 829–839. <https://doi.org/10.1016/j.bbagr.2010.09.004>

Acknowledgement

- 1) **Dr. K.V. Swamy**, Associate professor , Ph.D. Guide .
- 2) **Dr. D. Y. Patil Biotechnology and Bioinformatics Institute**, Pune.
- 3) **Prof. J. K. Pal**, Director, Dr. D. Y. Patil Biotechnology and Bioinformatics Institute, Pune.
- 4) **Prof. N.N. Nawani**, Research Coordinator, Dr. D. Y. Patil Biotechnology and Bioinformatics Institute, Pune.
- 5) **Mr. Kiran Lokhande**, ICMR SRF Bioinformatics Research Laboratory DYPBBI, Pune.

SCHRÖDINGER®



THANK YOU ☺